NITRIDERM® ULTRA ORANGE™

Nitrile Exam Gloves Extra Protection Series 199

Orange - The universal symbol for caution. Ideal for EMS, police and fire, hazmat functions, spills, etc. This medium weight nitrile glove is extremely elastic and comfortable while providing the tensile strength required for these demanding procedures. Polymer interior for rapid donning. Textured finish for an excellent grip even in the presence of moisture.

Characteristics: Powder-Free, Non-Sterile, Orange color

SIZE	ITEM NO.
XS	199050
S	199100
M	199200
L	199300
XL	199350
XXL	199400
XXXL	199450

XS-XL: 100 Gloves/Box, 10 Boxes/Case



Manufactured in compliance with multiple international standards, including ASTM D6319, ASTM D5151, ASTM F1671, and ASTM D6978



Features:

- Textured Finish = an Improved Wet/Dry Grip
- Low Modulus = a Softer, More Confortable Fit
- Non-Latex = No Risk of Latex Allergens

XXL-XXXL: 80 Gloves/Box, 10 Boxes/Case

·				_	
Tested for use with Chemotherapy Drugs as per ASTM D6978 Standard Practice for Assessment of Medical Gloves to Permeation by Chemotherapy Drugs Minimum Breakthrough Detection Time in Minutes:					
Azacytidine (25.0 mg/ml)	>240		Ifosfamide (50.0 mg/ml)	>240	
Carboplatin (10.0 mg/ml)	>240		Irinotecan (20.0 mg/ml)	>240	
Carmustine (3.3 mg/ml)	12.9		Methotrexate (25.0 mg/ml)	>240	
Cisplatin (1.0 mg/ml)	>240		Mitomycin C (0.5 mg/ml)	>240	
Cyclophosphamide (20.0 mg/ml)	>240		Mitoxantrone (2.0 mg/ml)	>240	
Dacarbazine (10.0 mg/ml)	>240		Oncovin (1.0 mg/ml)	>240	
Docetaxel (10.0 mg/ml)	>240		Oxaliplatin (5.0 mg/ml)	>240	
Doxorubicin Hydrochloride (2.0 mg/ml)	>240		Paclitaxel (6.0 mg/ml)	>240	
Epirubicin (2.0 mg/ml)	>240		Thiotepa (10.0 mg/ml)	45.0	
Etoposide (20.0 mg/ml)	>240		Vincristine Sulfate (1.0 mg/ml)	>240	
Fluorouracil (50.0 mg/ml)	>240		Vinorelbine (10.0 mg/ml)	>240	
Gemcitabine (38.0 mg/ml)	>240	Also Tested:	Fentanyl Citrate (100 mcg/2ml)	>240	

Tensile Strength & Elongation	Before Aging	After Accelerated Aging
Tensile Strength (Mpa)	32	27
ASTM Requirement Min. (Mpa)	14	14
Elongation (%)	650	480
ASTM Requirement Min. (%)	500	400

Average Length	9.5 in ◆ 240 mm
Average Palm Thickness	3.5 mil ◆ 0.09 mm
Average Finger Thickness	4.3 mil ♦ 0.11 mm

Please note that Carmustine and Thiotepa have extremely low permeation times of 12.9 minutes and 45.0 minutes respectively.

Warning: Do not use with Carmustine. Warning: Do not use with Thiotepa.

Caution: In compliance with ASTM D6978, the testing conditions used are intended to approximate the worst-case conditions for clinical use. Testing was conducted on single layer glove material. It is the users' responsibility to determine the applicability of these gloves for their intended use with chemotherapy drugs.



Corporation is certified to ISO 13485:2016 QMS for

medical devices.



Product Solutions You Trust

View this product on our website:

